

Abstract Word Count: 248

Manuscript Word Count: 3443

Can Individuals with Alcohol Use Disorder Achieve and Sustain Non-Abstinent Recovery? Non-

Abstinent Outcomes 10 Years After Alcohol Use Disorder Treatment

Katie Witkiewitz, PhD, Adam D. Wilson, MS

University of New Mexico

Corey R. Roos, PhD

Yale University

Julia E. Swan, BA, Victoria R. Votaw, MS, Elena R. Stein, MS, Matthew R. Pearson, PhD,

Karlyn A. Edwards, MS, J. Scott Tonigan, PhD

University of New Mexico

Kevin A. Hallgren, PhD

University of Washington

Kevin S. Montes, PhD

California State University Dominguez Hills

Stephen A. Maisto, PhD

Syracuse University

Jalie A. Tucker, PhD, MPH

University of Florida

Preparation of this manuscript was supported in part by grants from the National Institute on Alcohol Abuse and Alcoholism (R01 AA022328, 2K05 AA016928, K01 AA024796, K01 AA023233, and T32 AA018108). The authors declare no conflicts of interest.

Corresponding author: Katie Witkiewitz, University of New Mexico, MSC 03-2220,

Albuquerque NM 87131, (505) 925-2334, Fax: (505) 925-2301, katiw@unm.edu

Abstract

Objective: Four decades ago the “controlled drinking” controversy roiled the alcohol field. Data have subsequently accumulated indicating that non-abstinent alcohol use disorder (AUD) recovery is achievable, but questions remain whether it is sustainable long-term. This study examined whether non-abstinent recovery at three years following AUD treatment is associated with better functioning at 10 years following treatment. **Methods:** Data were from the 10-year follow-up of Project MATCH (New Mexico site only, $n=146$; 30.1% female, 58.6% non-White). Recovery was defined by latent profile analyses based on psychosocial functioning and alcohol consumption three years following treatment. Drinking practices and consequences, depression, purpose in life, and anger were assessed 10 years following treatment. Distal outcome analyses examined differences in drinking and functional outcomes at 10 years as a function of the 3-year latent profiles. Analyses were pre-registered at <https://osf.io/3hbxr>. **Results:** Four latent profiles identified at three years following treatment (i.e., low functioning frequent heavy drinkers, low functioning infrequent heavy drinkers, high functioning heavy drinkers, and high functioning infrequent non-heavy drinkers) were significantly associated with outcomes ten years following treatment. The two high functioning profiles at three years had the highest level of functioning at ten years post-treatment, regardless of alcohol consumption level. Abstinence at three years did not predict better functioning at ten years. **Conclusions:** Non-abstinent AUD recovery is possible and is sustainable for up to 10 years following treatment. The current findings align with recent proposals to move beyond relying on alcohol consumption as a central defining feature of AUD recovery.

Key Words: alcohol use disorder, alcohol treatment, recovery, mixture models, Project MATCH

Introduction

Abstinence from alcohol remains a primary target in many treatment programs for alcohol use disorder (AUD) ¹⁻³ and in mutual help organizations such as Alcoholics Anonymous (AA). Achieving abstinence is associated with improvements in psychosocial functioning ^{4,5} and lower risk of physical health problems ⁶⁻⁸. However, some individuals also attempt to recover from AUD by reducing their alcohol use (i.e., moderation or controlled drinking), and a number of empirical studies demonstrate that reducing alcohol consumption is also associated with improvements in functioning and health ⁹⁻¹².

Recent conceptualizations of the term ‘recovery’ have shifted to emphasize the broader biopsychosocial process of improvement that is related to, but not solely determined by, alcohol consumption. As noted by Ashford and colleagues¹³, stakeholder institutions, including the American Society of Addiction Medicine (ASAM), have updated and modified their operational definitions of recovery over the past two decades. In 2005, ASAM’s public policy statement on recovery highlighted a state of psychological and physical health in which an individual’s abstinence from substance use was “complete and comfortable”¹⁴. A 2013 update to ASAM’s definition still mentions “an individual’s consistent pursuit of abstinence,” but does so in the context of “a process of sustained action that addresses biological, psychological, social, and spiritual disturbances,” including improvements in interpersonal relationships, behavioral control, and emotional responding ¹⁵. Empirical support for a broader conceptualization of recovery has been reviewed in recent work ¹⁶⁻¹⁸ and also expressed by individuals who self-identify as being in recovery ¹⁹⁻²¹.

Consistent with expanding definitions of recovery, we recently conducted empirical investigations seeking to identify “recovery profiles” (i.e., subgroups) characterized by a range

of both alcohol-related and psychosocial functioning measures. Using data from the outpatient arm of Project MATCH²², we found four profiles at three years following AUD treatment: (1) low functioning frequent heavy drinkers (15.8% of the sample), (2) low functioning infrequent heavy drinkers (16.1%), (3) high functioning heavy drinkers (16.9%), and (4) high functioning, infrequent, non-heavy drinkers (51.2%)²³. These results suggest that abstinence may be a necessary recovery element for some individuals with AUD, but is not essential for all, and positive changes in functioning and well-being often are more fundamental elements.

Although this research adds to growing evidence that distinct longer term recovery profiles can be identified based on both alcohol-related outcomes and functioning indicators, important questions remain about whether these profiles forecast sustained positive outcomes over longer intervals. For example, do the individuals in the high functioning profiles—including those engaging in heavy drinking—maintain this level of functioning in subsequent years? A limited number of studies with follow-up intervals longer than three years suggest this is the case. For example, using follow-up data from the COMBINE Study²⁴, we found that individuals who were high functioning at three years post-treatment, regardless of level of alcohol consumption, had the best self-reported health and fewer hospitalizations at 7-9 years post-treatment²⁵. An observational study of individuals with AUD surveyed participants about their drinking practices, psychosocial functioning, and life contexts at baseline and 1, 3, 8, and 16 years later. Regardless of whether they had recently sought help or achieved abstinence, many participants showed improvement in alcohol-related functioning, life contexts, and coping²⁶. Taken together, these studies may inform a longstanding debate in the field concerning the risks and stability of non-abstinent recovery⁹ and the utility of broader conceptualizations of recovery that emphasize improvements in biopsychosocial functioning^{16,17}.

Current Study

The present study attempts to replicate and extend our recent work with the COMBINE Study²⁴ by investigating the relationships between recovery profiles identified at three years following treatment and alcohol consumption and functioning outcomes at ten years following treatment among a site-specific subset of participants involved in Project MATCH²².

Importantly, this is an extension of our analyses with COMBINE to include a broader set of psychosocial functioning indicators, for a longer time period, and the analyses are conducted in a more diverse sample of participants. We hypothesized the recovery typologies characterized by high functioning heavy drinking and high functioning infrequent non-heavy drinking at three years following treatment²³ would be associated with less drinking and higher psychosocial functioning at ten years following treatment when compared to recovery typologies characterized by low functioning frequent or infrequent heavy drinkers at three years following treatment.

Method

Participants and Procedure

The current study was a secondary data analysis of Project MATCH²², a randomized clinical trial comparing three psychosocial treatments for AUD: Cognitive Behavioral Therapy (CBT)²⁷, Motivational Enhancement Therapy (MET)²⁸, and Twelve-Step Facilitation (TSF)²⁹. Participants who sought outpatient treatment from nine research centers in the United States ($N=952$) and provided drinking data at the 3-year follow-up assessment ($n=806$; 84.7% retention) were included in the latent profile analysis. Additionally, participants in the outpatient arm at the Albuquerque, New Mexico site ($n=226$) were invited to complete a 10-year follow-up assessment³⁰. Of this subset, 149 provided data at the 10-year follow-up (66.4%) and 146 (64.6%) completed both the 3- and 10-year follow-up and were included in the present study.

Table 1 provides descriptive statistics for the subset of participants who provided 3-year and 10-year follow-up data.

Measures

Measures were assessed at multiple time points including baseline, three years following treatment (39-months post-baseline), and ten years following treatment.

Three-Year Follow-Up Latent Profile Indicators

Alcohol and drug use were assessed with the Form-90³¹, a calendar-based tool determining alcohol and drug use in the previous 90 days. Summary alcohol use variables included percent drinking days (PDD), percent heavy drinking days (PHDD, i.e., 4+ drinks/day for women, 5+ drinks/day for men), and drinks per drinking day (DDD). Marijuana and other illicit drug use responses were summarized as binary (0=no use, 1=any use).

The Drinker Inventory of Consequences (DrInC)³² was utilized to assess negative alcohol-related consequences. Participants reported the regularity of 45 consequences on a 4-point scale (1=never, 4=daily or almost daily). Internal consistency of the DrInC for this sample was $\alpha=0.97$.

The Psychosocial Functioning Inventory (PFI)³³ was used to evaluate social functioning. The 10-item social behavior subscale from the PFI included items assessing the prevalence of problem social interactions and behaviors in the past 30 days (e.g., “Demanded others do things your way”); higher scores indicated better psychosocial functioning. Sample internal consistency of the PFI social behavior subscale was $\alpha=0.83$. Additionally, we selected four items to measure life satisfaction and social functioning over the past 30 days with binary responses (0=satisfied/happy; 1=dissatisfied/unhappy). The internal consistency of these four items was $\alpha=0.79$.

The Addiction Severity Index (ASI)³⁴ was used to measure employment and experiences with affective or cognitive symptoms (e.g., “severe depression”) in the past 30 days with responses coded as binary (0=employed or symptom not present, 1=unemployed or symptom present). We examined four individual items given prior evidence of poor measurement properties of the ASI composites³⁵. The internal consistency of the four items was $\alpha=0.63$.

10-Year Follow-Up Outcomes

Different measures from the original Project MATCH study were assessed at the 10-year follow-up assessment, and only the Form-90 and DrInC were administered at both 3-year and 10-year follow-ups. The PFI and ASI were not administered at the 10-year follow-up.

The Form-90³⁶ was utilized at the 10-year follow-up to assess alcohol use, including PDD, PHDD, and DDD. These variables represent alcohol use over the 30 days prior to the 10-year assessment.

The DrInC³² was also used to measure the frequency of 45 alcohol-related negative consequences over the three months prior to the 10-year follow-up assessment. At this assessment, internal consistency of the DrInC was $\alpha=0.98$.

The Beck Depression Inventory (BDI)³⁷ was utilized to assess 21 depression symptoms (rated from 0 to 3, representing the severity of each symptom) in the previous week. At the 10-year follow-up, the BDI demonstrated internal consistency of $\alpha=0.91$.

The Purpose in Life (PIL) test³⁸ was used to measure individuals’ sense of meaning or purpose in life, including items assessing life meaning, satisfaction, freedom, fear of death, suicidal ideation, and how worthwhile one’s life is. This 20-item questionnaire had an internal consistency of $\alpha=0.92$ at the 10-year assessment.

The State-Trait Anger Expression Inventory (STAXI) assessed the expression of anger in

situational contexts (state anger) and a predisposition toward anger (trait anger)³⁹. The total score from the 44-item questionnaire was utilized in the present analysis, and the internal consistency was $\alpha=0.87$.

Analytic Approach

Analyses for the current study were preregistered. Project documents and preregistration details can be found at <https://osf.io/3hbxx> (last accessed 6/16/2020). Latent profiles of drinking and functioning at the 3-year follow-up were derived using latent profile analysis in *Mplus* version 8⁴⁰. The latent profile models were estimated based on the entire Project MATCH outpatient sample who completed the 3-year follow-up assessment ($n=806$; 84.6% of the MATCH outpatient sample). As detailed by Witkiewitz et al.²³, we used a weighted maximum likelihood function and estimated variance-covariance matrix for all available 3-year data ($n=806$). Model fit was examined using the Lo Mendell Rubin Likelihood Ratio test (LRT), Bayesian Information Criterion (BIC), and sample-size-adjusted BIC (aBIC). Classification precision was defined using relative entropy, where entropy greater than .80 indicated good classification precision⁴¹.

Distal Outcomes at 10-Year Follow-Up

To examine differences in latent profiles on functioning at 10 years following treatment, we conducted latent profile analyses with the manual 3-step Bolck, Croons, and Hageaars (BCH) approach⁴²⁻⁴⁴ to assess whether the latent profiles at the 3-year follow-up were associated with distal outcomes of alcohol consumption, alcohol consequences, and psychosocial functioning at the 10-year follow-up. Of the Albuquerque outpatient sample that was consented to the 10-year follow-up ($n=193$), 149 had data at the 10-year follow-up (66.4%) and 146 (64.6%) completed both the 3- and 10-year follow-up, and were included in the distal outcomes

analysis. Differences in distal outcomes at 10 years as a function of latent profile membership at 3 years were evaluated using a Wald χ^2 test with paired *t*-tests for pairwise profile comparisons. Cohen's *d* statistic, which reflects standardized mean differences in outcomes (i.e., differences between groups in standard deviation units), was also calculated. A priori power analyses, using a Monte Carlo simulation, indicated power to detect distal outcome effects (i.e., mean differences in 10-year outcomes) by profile membership at 10 years was greater than 0.63 to detect medium effect sizes and we had power greater than 0.97 to detect large effect sizes.

Sensitivity Analyses. Attrition analyses (*t*-tests for continuous measures and χ^2 tests for categorical measures) were used to explore differences in the baseline and 3-year follow-up assessments among those with complete data at the 10-year follow-up ($n=149$) versus those who did not complete the 10-year follow-up ($n=43$). We also examined whether there were differences in latent profiles at the 3-year assessment and completion of the 10-year follow-up assessment. Additional analyses were conducted, which were not preregistered, to examine differences in 10-year outcomes among those who were or were not abstinent at the 3-year follow-up. Mean differences were examined using Wald χ^2 tests with *t*-tests for individual outcome comparisons. Cohen's *d* statistics were also calculated comparing abstainers to non-abstainers.

Results

Latent Profiles at the 3-Year Follow-Up

As reported previously²³, the four-profile model at the three-year follow-up was selected based on BIC, aBIC, and LRT statistics, and the model classification precision was excellent (entropy=0.92). The profiles were substantively meaningful and were labeled “low functioning frequent heavy drinking” (profile 1; 15.8% of the sample), “low functioning infrequent heavy

drinking” (profile 2; 16.1%, “high functioning heavy drinking” (profile 3; 16.9%), and “high functioning infrequent non-heavy drinking” (profile 4; 51.2%). None of the individuals with expected classification in profile 1 (low functioning frequent heavy drinking) or profile 3 (high functioning heavy drinking) were abstinent. Of those with expected classification in profile 4 (high functioning infrequent non-heavy drinking), 48.9% were abstinent from alcohol; and of those with expected classification in profile 2 (low functioning infrequent heavy drinking), 26.9% were abstinent from alcohol.

Distal Outcomes at the 10-Year Follow-Up

Table 1 provides descriptive statistics for distal outcomes by latent profile membership, which differed significantly at the 10-year follow-up (Wald $\chi^2(24)=133.85, p < 0.001$). Figure 1 depicts alcohol consumption and alcohol-related consequences at the 10-year follow-up among individuals with expected classification in profiles 1-4, respectively. Pairwise comparisons indicated that those in profile 4 (high functioning infrequent non-heavy drinking) had significantly fewer consequences on the DrInC, as compared to all other profiles (all $p < 0.04$), and had significantly fewer drinks per drinking day ($t(1)=-4.25, p=0.04$), percent drinking days ($t(1)=-36.37, p=0.001$), and percent heavy drinking days ($t(1)=-17.82, p=0.04$), as compared to profile 3 (high functioning heavy drinking). Profile 3 also reported significantly greater percent drinking days, as compared to profile 1 ($t(1)=31.76, p=0.02$) and profile 2 ($t(1)=29.19, p=0.02$). We also examined abstinence status at the 10-year follow-up by latent profile membership at 3 years and found that 62.5% of individuals in profile 1, 39.1% of individuals in profile 2, 19.0% of individuals in profile 3, and 50.0% of individuals in profile 4 were abstinent in the month preceding the 10-year follow-up assessment.

Functional outcomes at 10-year follow-up (Figure 2), indicated profiles 3 and 4 had

significantly greater purpose in life, as compared to profile 2 (all $ps < .002$). Profile 4 had significantly lower depression than profile 1 ($t(1)=-5.53, p=0.007$) and profile 2 ($t(1)=-9.22, p < 0.001$), but did not significantly differ from profile 3 ($t(1)=-4.31, p=0.06$). Profile 4 also had significantly less anger than profile 3 ($t(1)=-8.62, p=0.02$) and profile 2 ($t(1)=-16.69, p < 0.001$), and profile 2 had significantly more anger than profile 1 ($t(1) =14.21, p= 0.01$).

Sensitivity Analyses

Attrition analyses indicated no significant differences on any study variables (alcohol consumption, consequences, or psychosocial functioning) at baseline or at the 3-year follow-up between those who did ($n=149$) and did not complete ($n=43$) the 10-year follow-up. Of the 806 outpatient participants who had 3-year data used to determine their latent profile, a χ^2 test of independence showed that profile membership of the 10-year retained sample did not significantly differ from those lost to follow-up ($n=660; \chi^2(3)=4.66, p=.199$), nor did profile membership differ from the Albuquerque subsample who were eligible but did not complete the 10-year follow-up ($n=47; \chi^2(3)=1.554, p=.670$).

Additional analyses were conducted to examine mean outcome differences at the 10-year follow-up among those who were ($n=33$) and were not abstinent ($n=113$) at the 3-year follow-up. Significant differences in distal outcomes were found between groups at the 10-year follow-up (Wald $\chi^2(7)=20.79, p=0.004$). Supplementary Figures 1 and 2 illustrate the differences in mean outcomes (with standard errors indicated by error bars) and Cohen's d standardized mean differences between abstainers and drinkers. Abstainers at the 3-year follow-up reported significantly lower percent heavy drinking days ($t(1)=-11.04, p=0.01$), percent drinking days ($t(1)=-19.02, p < 0.001$), and drinking consequences ($t(1) =-12.68, p=0.02$) at the 10-year follow-up, but did not differ on any of the other functional outcomes (all $ps > .13$).

Discussion

As hypothesized, the two highest functioning profiles at three years following treatment (profile 3 and 4) generally had the best functioning outcomes, including greater purpose in life and lower levels of depression, at ten years following treatment. These findings support the clinical validity of the recovery profiles and reaffirm the importance of considering indicators of functioning, and not simply alcohol consumption levels, when defining long-term recovery from AUD. Indeed, our findings revealed a lack of a one-to-one correspondence between drinking behavior and functioning during the process of recovery over time. Abstinence three years following treatment did not predict better functioning ten years following treatment. Rather, *functioning* at three years following treatment (profiles 3 and 4) predicted better *functioning* at ten years following treatment. As recently proposed, focusing on functioning rather than drinking practices per se may be more useful when defining successful AUD recovery and forecasting how an individual will fare over the long run^{4,13,167,17,45}. Our quantitative findings using a clinical treatment sample also are aligned with research indicating that functional outcomes, including quality of life and well-being, are highly valued among persons who self-identify as being in recovery^{20,21}.

An important finding was that profile 3, which was characterized by occasional heavy drinking three years following treatment, generally showed favorable functioning outcomes at both the three-year and ten-year follow-ups. For example, compared to low functioning profiles 1 and 2, membership in profile 3 was associated with significantly higher purpose in life at year 10, whereas depression symptoms and purpose in life were similar in high functioning profiles 3 and 4. These findings are consistent with our prior work demonstrating that heavy drinking itself may not be a valid outcome indicator among individuals with AUD^{4,45-47}. Overall, study

findings support and extend a growing body of literature that suggests that abstinence is not the only path to achieve and/or sustain a high level of psychosocial functioning in the years following alcohol treatment. Rather, there appear to be multiple paths to recovery that can include moderate or heavy alcohol consumption post-treatment for some individuals^{16,23,25,48}.

It is also important to note that profile 4 (high functioning, infrequent non-heavy drinking at year 3) had the best overall outcomes at year 10, despite the fact that some individuals in profile 4 had returned to some heavy drinking (average percent heavy drinking days of 10%) at year 10. Profile 4 had significantly lower anger, depression, and alcohol-related consequences, and greater purpose in life than profiles 1 and 2 and did not differ significantly from profile 3 on two important functioning outcomes at year 10 (i.e., depression and purpose in life). While individuals who achieved both high functioning and abstinence/non-heavy drinking (profile 4) at three years had optimal long-term outcomes as a whole, individuals who have a combination of high functioning and more frequent heavy drinking (profile 3) also showed favorable long-term outcomes in psychosocial functioning.

Some differences in ten-year drinking outcomes among recovery profiles at three years following treatment were somewhat unexpected. For instance, rates of abstinence at year 10 for profiles 1, 2, 3, and 4 were 62.5%, 39.1%, 19.0%, and 50.0%, respectively. It is quite possible that individuals in profile 1 (low functioning frequent heavy drinking at year 3) were not able to sustain frequent heavy drinking without significant negative consequences and thus were more likely to abstain at 10 years. Additionally, those in high functioning profile 3 reported the highest levels of drinks per drinking day, percent drinking days, and percent heavy drinking days at year 10 – but lower levels of drinking consequences at year 10 – relative to the two low functioning profiles (profiles 1 and 2).

Limitations and Future Directions

Several limitations should be considered when interpreting the present results. First, the current study examined profiles of recovery and long-term outcomes among an outpatient sample collected at one site over a decade ago, and findings may not generalize to inpatient or more contemporary samples. Replication in larger samples of individuals who received inpatient and outpatient treatment is warranted. Second, recovery is likely best conceptualized as a dynamic process of change¹⁷, and the present study only examined outcomes in a three-month timeframe ten years following AUD treatment. It did not assess behavior patterns and outcomes or capture other treatment that may have occurred during the 10 years following the Project MATCH treatment episode, and the extent of additional help-seeking across time is unknown. Third, all measures included in the study were self-reported, and collateral information from providers or family were not available. We also did not measure physical health or medical outcomes at year 10, and we were not able to evaluate whether individuals who continued drinking heavily were at increased risk for physical health-related diseases⁴⁸⁻⁵⁰. Future studies should include additional indicators and outcomes, such as cognitive functioning, chronic pain, medical comorbidities, and other substance use to advance our understanding of functioning following AUD treatment. Fourth, only a small subset of original participants in the outpatient arm of Project MATCH at the Albuquerque site were followed successfully over 10 years, and the extent to which the sample may be biased cannot be thoroughly ascertained. Nevertheless, alcohol consumption, consequences, and psychosocial functioning at baseline and at the 3-year follow-up did not differ significantly between participants who did ($n=149$) and did not complete ($n=43$) the 10-year follow-up.

With these qualifications, the present study adds to evidence that non-abstinent AUD

recovery is possible and can be maintained for up to 10 years following treatment. The findings support recent proposals to move beyond viewing abstinence as a central defining feature of AUD recovery and relying heavily on quantity-frequency measures of drinking practices as the primary outcome indicator. Future research that expands the scope of outcome indicators to include measures of biopsychosocial functioning is important for advancing understanding of the multiple pathways to recovery from AUD.

References

1. Davis AK, Rosenberg H, Rosansky JA. American counselors' acceptance of non-abstinence outcome goals for clients diagnosed with co-occurring substance use and other psychiatric disorders. *J Subst Abuse Treat.* 2017;82:29-33. doi:10.1016/j.jsat.2017.08.006
2. Rosenberg H, Davis LA. Acceptance of moderate drinking by alcohol treatment services in the United States. *J Stud Alcohol.* 1994;55(2):167-172.
<http://www.ncbi.nlm.nih.gov/pubmed/8189737>. Accessed July 5, 2013.
3. Davis AK, Rosenberg H. Acceptance of non-abstinence goals by addiction professionals in the United States.: Bulletin of the society of psychologists in addictive behaviors: Bulletin of the society of psychologists in substance abuse. *Psychol Addict Behav.* 2013;27(4):1102-1109. doi:10.1037/a0030563
4. Wilson AD, Bravo AJ, Pearson MR, Witkiewitz K. Finding success in failure: using latent profile analysis to examine heterogeneity in psychosocial functioning among heavy drinkers following treatment. *Addiction.* 2016;111(12):2145-2154. doi:10.1111/add.13518
5. Maisto SA, Clifford P r., Longabauch R, Beattie M. The relationship between abstinence for one year following pretreatment assessment and alcohol use and other functioning at two years in individuals presenting for alcohol treatment. *J Stud Alcohol.* 2002;July:397-403.
6. GBD 2016 Alcohol Collaborators MG, Fullman N, Hawley C, et al. Alcohol use and burden for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet.* 2018;392(10152):1015-1035. doi:10.1016/S0140-6736(18)31310-2
7. Zemore SE, Lui C, Mericle A, Hemberg J, Kaskutas LA. A longitudinal study of the

- comparative efficacy of Women for Sobriety, LifeRing, SMART Recovery, and 12-step groups for those with AUD. *J Subst Abuse Treat.* 2018;88:18-26.
doi:10.1016/j.jsat.2018.02.004
8. de Visser RO, Piper R. Short- and Longer-Term Benefits of Temporary Alcohol Abstinence During ‘Dry January’ Are Not Also Observed Among Adult Drinkers in the General Population: Prospective Cohort Study. *Alcohol Alcohol.* 2020.
doi:10.1093/alcalc/aaa025
 9. Sobell MB, Sobell LC. The aftermath of heresy: A response to Pendery et al.’s (1982) critique of “Individualized behavior therapy for alcoholics.” *Behav Res Ther.* 1984;22(4):413-440. doi:10.1016/0005-7967(84)90084-6
 10. Sobell LC, Cunningham JA, Sobell MB. Recovery from alcohol problems with and without treatment: Prevalence in two population surveys. *Am J Public Health.* 1996;86(7):966-972. doi:10.2105/AJPH.86.7.966
 11. Cheong JW, Lindstrom K, Chandler SD, Tucker JA. Utility of different dimensional properties of drinking practices to predict stable low-risk drinking outcomes of natural recovery attempts. *Addict Behav.* 2020;106:106387. doi:10.1016/j.addbeh.2020.106387
 12. Fan AZ, Chou SP, Zhang H, Jung J, Grant BF. Prevalence and correlates of past year recovery from DSM-5 alcohol use disorder: Results from National Epidemiologic Survey on Alcohol and Related Conditions-III. *Alcohol Clin Exp Res.* 2019;43(11):2406-2420.
 13. Ashford RD, Brown A, Brown T, et al. Defining and operationalizing the phenomena of recovery: a working definition from the recovery science research collaborative. *Addict Res Theory.* 2019;27(3):179-188. doi:10.1080/16066359.2018.1515352
 14. American Society of Addiction Medicine. *Public Policy Statement on the State of*

- Recovery*. Chevy Chase, MD, USA; 2005.
15. American Society of Addiction Medicine. *Terminology Related to Addiction, Treatment, and Recovery*. Chevy Chase, MD, USA; 2013.
 16. Witkiewitz K, Montes KS, Schwebel FJ, Tucker JA. What is recovery? A narrative review of definitions of recovery from alcohol use disorder. *Alcohol Res Curr Rev*.
 17. Witkiewitz K, Tucker JA. Abstinence not required: Expanding the definition of recovery from alcohol use disorder. *Alcohol Clin Exp Res*. 2020;44(1):36-40.
doi:10.1111/acer.14235
 18. Neale J, Finch E, Marsden J, et al. How should we measure addiction recovery? Analysis of service provider perspectives using online Delphi groups. *Drugs Educ Prev Policy*. 2014;21(4):310-323. doi:10.3109/09687637.2014.918089
 19. Kaskutas LA, Borkman TJ, Laudet A, et al. Elements that define recovery: the experiential perspective. *J Stud Alcohol Drugs*. 2014;75(6):999-1010.
doi:10.15288/jsad.2014.75.999
 20. Kelly JF, Greene MC, Bergman BG. Beyond abstinence: Changes in indices of quality of life with time in recovery in a nationally representative sample of U.S. adults. *Alcohol Clin Exp Res*. 2018;42(4):770-780. doi:10.1111/acer.13604
 21. Neale J, Vitoratou S, Finch E, et al. Development and validation of SURE: A patient reported outcome measure for recovery from drug and alcohol dependence. *Drug Alcohol Depend*. 2016;165:159-167. doi:10.1016/j.drugalcdep.2016.06.006
 22. Project MATCH Research Group. Matching Alcoholism Treatments to Client Heterogeneity: Project MATCH posttreatment drinking outcomes. *J Stud Alcohol*. 1997;58(1):7-29. <http://www.ncbi.nlm.nih.gov/pubmed/8979210>. Accessed September 27,

- 2013.
23. Witkiewitz K, Wilson AD, Pearson MR, et al. Profiles of recovery from alcohol use disorder at three years following treatment: Can the definition of recovery be extended to include high functioning heavy drinkers? *Addiction*. 2019;114(1):69-80. doi:10.1111/add.14403
 24. Anton RF, O'Malley SS, Ciraulo DA, et al. Combined pharmacotherapies and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*. 2006;295(17):2003-2017. doi:10.1001/jama.295.17.2003
 25. Witkiewitz K, Pearson MR, Wilson AD, et al. Can alcohol use disorder recovery include some heavy drinking? A replication and extension up to nine years following treatment.
 26. Moos RH, Moos BS. Sixteen-year changes and stable remission among treated and untreated individuals with alcohol use disorders. *Drug Alcohol Depend*. 2005;80(3):337-347. doi:10.1016/j.drugalcdep.2005.05.001
 27. Kadden RM, Carroll KM, Donovan DM, et al. *Cognitive-Behavioral Coping Skills Therapy Manual*. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; 1995. [https://pubs.niaaa.nih.gov/publications/MATCHSeries3/Project MATCH Vol_3.pdf](https://pubs.niaaa.nih.gov/publications/MATCHSeries3/Project%20MATCH%20Vol_3.pdf).
 28. Miller WR, Zweben A, DiClemente CC, Rychtarik RG. *Motivation Enhancement Therapy Manual*. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; 1994.
 29. Nowinski J, Baker S, Carroll KM. *Twelve Step Facilitation Therapy Manual*. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism; 1995.
 30. Pagano ME, White WL, Kelly JF, Stout RL, Tonigan JS. The 10-year course of Alcoholics Anonymous participation and long-term outcomes: a follow-up study of

- outpatient subjects in Project MATCH. *Subst Abus.* 2013;34(1):51-59.
doi:10.1080/08897077.2012.691450
31. Miller WR, National Institute on Alcohol Abuse and Alcoholism. *Form 90: A Structured Assessment Interview for Drinking and Related Behaviors (Test Manual)*, NIAAA Project MATCH Monograph Series, Vol. 5.; 1996.
 32. Miller WR, Tonigan JS, Longabaugh R. The Drinker Inventory of Consequences (DrInC). An Instrument for Assessing Adverse Consequences of Alcohol Abuse. Test Manual. ... *MATCH Monogr Ser.* 1995.
 33. Feragne MA, Longabaugh R, Stevenson JF. The psychosocial functioning inventory. *Eval Health Prof.* 1983;6(1):25-48. doi:10.1177/016327878300600102
 34. McLellan AT, Kushner H, Metzger D, et al. The fifth edition of the addiction severity index. *J Subst Abuse Treat.* 1992;9(3):199-213. doi:10.1016/0740-5472(92)90062-S
 35. Mäkelä K. Studies of the reliability and validity of the Addiction Severity Index. *Addiction.* 2004;99(4):398-410. doi:10.1111/j.1360-0443.2003.00665.x
 36. Miller WR. *Form 90: A Structured Assessment Interview for Drinking and Related Behaviors*. Project MA. Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism; 1996.
 37. Beck AT, Steer RA, Carbin MG. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clin Psychol Rev.* 1988;8(1):77-100.
doi:10.1016/0272-7358(88)90050-5
 38. Crumbaugh JC, Maholick LT. An experimental study in existentialism: The psychometric approach to Frankl's concept of noogenic neurosis. *J Clin Psychol.* 1964;20(2):200-207.
doi:10.1002/1097-4679(196404)20:2<200::AID-JCLP2270200203>3.0.CO;2-U

39. Spielberger CD, Sydeman SJ. State-trait anxiety inventory and state-trait anger expression inventory. In: *The Use of Psychological Testing for Treatment Planning and Outcome Assessment.* ; 1994.
40. Muthén LK, Muthén BO. Mplus users guide (Version 8). 2019.
41. Nylund KL, Asparouhov T, Muthén BO. Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Struct Equ Model.* 2007;14(4):535-569.
42. Bolck A, Croon M, Hagenaars JA. Estimating latent structure models with categorical variables: One-step versus three-step estimators. *Polit Anal.* 2004;12(1):3-27.
doi:10.1093/pan/mph001
43. Nylund-Gibson K, Grimm RP, Masyn KE. Prediction from latent classes: A demonstration of different approaches to include distal outcomes in mixture models. *Struct Equ Model A Multidiscip J.* 2019;26(6):967-985.
doi:10.1080/10705511.2019.1590146
44. Yeo RA, Phillips JP, Jung RE, Brown AJ, Campbell RC, Brooks WM. Magnetic resonance spectroscopy detects brain injury and predicts cognitive functioning in children with brain injuries. *J Neurotrauma.* 2006;23(10):1427-1435.
doi:10.1089/neu.2006.23.1427
45. Maisto SA, Hallgren KA, Roos CR, Witkiewitz K. Course of remission from and relapse to heavy drinking following outpatient treatment of alcohol use disorder. *Drug Alcohol Depend.* 2018;187:319-326. doi:10.1016/j.drugalcdep.2018.03.011
46. Tucker JA, Cheong J, James T, Jung S, Chandler SD. Pre-resolution drinking problem severity profiles associated with stable moderation outcomes of natural recovery attempts.

Alcohol Clin Exp Res. 2020;44(3):738-748. doi:10.1111/acer.14287


47. Witkiewitz K, Kirouac M, Roos CR, et al. Abstinence and low risk drinking during treatment: Association with psychosocial functioning, alcohol use, and alcohol problems 3 years following treatment. *Psychol Addict Behav.* 2018;32(6):639-646.
doi:10.1037/adb0000381
48. Tucker JA, Chandler SD, Witkiewitz K. Epidemiology of recovery from alcohol use disorder. *Alcohol Res Curr Rev.* 2020.

Table 1. *Descriptive Statistics for 10-Year Sample by 3-Year Latent Profile Membership (Analysis N=146)*

	Total Sample (N=146)	Profile 1: Low Functioning Frequent Heavy Drinking (n=16)	Profile 2: Low Functioning Infrequent Heavy Drinking (n=23)	Profile 3: High Functioning Heavy Drinking (n=22)	Profile 4: High Functioning Infrequent Drinking (n=85)
10-Year Outcomes	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)
Percent drinking days	24.83 (33.98)	23.33 (36.64)	25.07 (31.62)	50.95 (42.51)	18.45 (28.76)
Percent heavy drinking days	14.85 (27.81)	22.71 (36.23)	17.25 (28.33)	25.71 (35.31)	9.60 (22.37)
Drinks per drinking day	5.04 (7.81)	5.33 (10.55)	6.71 (8.13)	7.66 (8.00)	3.86 (6.89)
DrInC Total Score	24.21 (30.07)	36.69 (37.77)	41.67 (38.80)	29.52 (25.65)	16.96 (24.57)
Beck Depression Inventory	7.92 (8.02)	10.81 (7.61)	13.81 (8.32)	9.43 (9.00)	5.54 (6.80)
Purpose in Life	103.82 (16.67)	98.75 (15.17)	91.24 (15.47)	106.10 (16.50)	106.71 (15.47)
State-Trait Anger Expression Inventory	80.46 (13.30)	79.25 (13.40)	91.86 (18.64)	84.67 (14.42)	77.08 (9.43)
Demographics					
Age (Baseline)	42.83 (8.59)	43.38 (7.66)	43.23 (7.41)	44.00 (9.71)	42.32 (8.83)
Sex (N, % Male)	103(71.0)	11 (68.8)	12 (57.1)	14 (66.7)	63 (75.0)
Marital status (N, % Married)	45 (31.0)	4 (25.0)	6 (28.6)	7 (33.3)	28 (33.3)
Race (N, % White)	60 (41.4)	7 (43.8)	9 (42.9)	11 (52.4)	31 (36.9)


Note. DrInC=Drinker Inventory of Consequences. Sample size (n) for each profile is based on the most likely class membership.

Figure 1. *Drinking outcomes ten years following treatment as a function of latent profile membership three years following treatment*




Note. Standard errors indicated by error bars. Cohen’s d standardized mean differences shown between profiles using profile 1 (low functioning frequent heavy drinking) and profile 4 (high functioning infrequent drinking) as reference groups.

Figure 2. *Functioning outcomes ten years following treatment as a function of latent profile membership three years following treatment*




Note. Standard errors indicated by error bars. Cohen’s d standardized mean differences shown between profiles using profile 1 (low functioning frequent heavy drinking) and profile 4 (high functioning infrequent drinking) as reference groups.

Supplementary Figure 1. *Drinking outcomes ten years following treatment by abstainer status at three years following treatment*



Note. Standard errors indicated by error bars). Cohen’s d standardized mean differences shown by abstainer status at three years following treatment.

Supplementary Figure 2. *Functioning outcomes ten years following treatment by abstainer status at three years following treatment*

Note. Standard errors indicated by error bars). Cohen's d standardized mean differences shown by abstainer status at three years following treatment.